

# Mobile genetic elements and their contribution to the emergence of antimicrobial resistant *Enterococcus faecalis* and *Enterococcus faecium*

K. Hegstad<sup>1,2</sup>, T. Mikalsen<sup>2</sup>, T. M. Coque<sup>3</sup>, G. Werner<sup>4</sup> and A. Sundsfjord<sup>1,2</sup>

1) Reference Centre for Detection of Antimicrobial Resistance, Department of Microbiology and Infection Control, University Hospital of North-Norway and  
2) Research group for Host-Microbe Interactions, Department of Medical Biology, University of Tromsø, Tromsø, Norway, 3) University Hospital Ramón y Cajal; Unidad de Resistencia a Antibióticos y Virulencia Bacteriana (RYC-CSIC), CIBER en Epidemiología y Salud Pública (CIBER-ESP), Madrid, Spain and  
4) Robert Koch-Institute, Wernigerode Branch, Wernigerode, Germany

## Abstract

Mobile genetic elements (MGEs) including plasmids and transposons are pivotal in the dissemination and persistence of antimicrobial resistance in *Enterococcus faecalis* and *Enterococcus faecium*. Enterococcal MGEs have also been shown to be able to transfer resistance determinants to more pathogenic bacteria such as *Staphylococcus aureus*. Despite their importance, we have a limited knowledge about the prevalence, distribution and genetic content of specific MGEs in enterococcal populations. Molecular epidemiological studies of enterococcal MGEs have been hampered by the lack of standardized molecular typing methods and relevant genome information. This review focuses on recent developments in the detection of MGEs and their contribution to the spread of antimicrobial resistance in clinically relevant enterococci.

**Keywords:** antimicrobial resistance, enterococcus, insertion sequence, mobile genetic elements, plasmid, review, transposon

*Clin Microbiol Infect* 2010; **16**: 541–554

**Corresponding author and reprint requests:** K. Hegstad, Reference Centre for Detection of Antimicrobial Resistance, Department of Microbiology and Infection Control, University Hospital of North-Norway, N-9038 Tromsø, Norway  
**E-mail:** Kristin.Hegstad@uit.no

## Introduction

Enterococci are important nosocomial pathogens [1]. They are uniquely armed for the antibiotic era and express intrinsic reduced susceptibility to major classes of antimicrobial agents and biocides [2,3]. The high propensity of enterococci to acquire and express new resistance determinants further enhances their ability to sustain antibiotic selection, promoting gastrointestinal colonization and nosocomial infections by antibiotic-resistant enterococci [4].

Transferable antimicrobial resistance in enterococci was first described in the early 1970s [5–7]. The detection and molecular clarification of transferable high-level vancomycin resistance in *Enterococcus faecium* in the late 1980s have further fuelled our interest in the mechanisms and routes to antimicrobial resistance in enterococci [8–11]. Importantly, as residents of human and animal bowel flora, enterococci are in a position to acquire resistance genes from other commensals, which may subsequently proceed to other more pathogenic bacteria [12,13].

Molecular biological studies have elucidated complex functional properties of important mobile genetics elements (MGEs) involved in transferable resistance in enterococci [10,14,15]. However, there is a considerable gap in our knowledge on the molecular epidemiology of MGEs, their genetic content and composition, as well as their relative contributions to the spread of defined antimicrobial resistances. Recent progress in molecular typing methods and enterococcal genome information has provided new tools and necessary insights for filling this gap. This review focuses on MGEs involved in the spread and expression of clinically important antimicrobial resistance in enterococci and their potential contribution to the spread of hospital-adapted clonal lineages of *E. faecium* and *Enterococcus faecalis*.

## Plasmids as Important Vehicles for Genetic Information in Enterococci

By definition, plasmids are semi-autonomously replicating extrachromosomal genetic elements. Differences in replica-

tion strategies and modular structures profoundly affect plasmid properties, such as size, copy number, host dependence and host range [16]. The essential backbones for successful plasmids include genetic modules encoding self-replication, stable inheritance and the ability to transfer between bacteria. Accessory plasmid content is integrated in between functional plasmid backbone modules and represents a huge reservoir of genetic variability, often with unknown functions, that is shared among different bacterial genera through horizontal gene transfer.

There are several criteria to classify plasmids in general and plasmids related to Gram-positive bacteria in particular. The mode of replication has been used to distinguish rolling circle replication plasmids and theta-replicating plasmids. In addition, plasmids that fail to co-reside in the same cell are grouped in incompatibility (Inc) groups [17,18]. Inc18 plasmids constitute a large group of enterococcal/streptococcal plasmids with a broad host range [19,20]. Pheromone-responsive plasmids represent a unique group of self-transferable (conjugative) narrow host range plasmids mostly described in *E. faecalis* [21]. pADI, pAM373 and pCF10 are well-known examples of pheromone-responsive plasmids, where the conjugative process is initiated as a response to short peptide pheromones produced by pheromone-responsive plasmid-free recipient strains mediating intercellular aggregation and high-frequency DNA transfer. Recently, Weaver *et al.* [22] proposed a new family (RepA<sub>N</sub>) of broadly distributed plasmids in Gram-positive bacteria encompassing pheromone-responsive plasmids of *E. faecalis*, as well as pRUM of *E. faecium*. Detailed sequence comparisons of the replication initiator protein suggest that the replicons have evolved along with their specific host, explaining their relatively narrow host range.

Plasmid replicon modules (replicons) have recently been used as targets of more simplistic methods for typing and epidemiological tracing of plasmids conferring antimicrobial resistance (R-plasmids). Other essential gene sets for plasmid survival, such as mobilization regions, have also been suggested as targets [23]. Given the modular evolution and genetic plasticity of plasmids, it is of note that schemes based on different core elements may not be congruent [16]. Carattoli *et al.* [18] developed a PCR-based plasmid typing method based on the replication regions from various plasmid incompatibility groups occurring in Enterobacteriaceae. A similar approach was recently described for the detection of plasmids from enterococci and other Gram-positive bacteria [24]. On the basis of 111 published sequences from Gram-positive bacteria, 19 replicon families (*rep*-family) and several unique replicons were identified. Using this PCR-based typing system, pCF10 (*rep*<sub>9</sub>)-, Inc18 (*rep*<sub>1</sub> and *rep*<sub>2</sub>)-

and pUSA02 (*rep*<sub>7</sub>)-related replicons were identified as being most prevalent in *E. faecalis* strains (*n* = 28), whereas Inc18 (*rep*<sub>2</sub>)-, pRII (*rep*<sub>14</sub>)- and pRUM (*rep*<sub>17</sub>)-related replicons dominated in *E. faecium* strains (*n* = 51) of human and animal origin [24]. However, approximately 30% of the strains tested did not support any *rep*-detection, indicating the presence of unidentified *rep*-types.

The enterococcal-specific parts of the *rep*-detection system described by Jensen *et al.* [24] have been used by others. A recent study of an epidemiologically diverse collection of *E. faecium* strains (*n* = 93) revealed a high prevalence of Inc18 (*rep*<sub>2</sub>)-, pRUM (*rep*<sub>17</sub>)- and pHTβ (*rep*<sub>unique</sub>)-related replicons [25]. The actual enterococcal typing scheme accounted for approximately 60% of the total number of plasmids visualized by S1-nuclease analyses. Interestingly, strains belonging to hospital-adapted clades (CC17-related) yielded a significant higher number of *rep* types and pRUM (*rep*<sub>17</sub>)-related replicons in particular, indicating a role in accessory plasmid DNA for promoting hospital adaptation. *Rep*-typing of extended *E. faecalis* strain collections has so far shown a dominance of pheromone-responsive plasmid (*rep*<sub>8</sub> and *9*)-, pS86 (*rep*<sub>6</sub>)- and Inc18 (*rep*<sub>1</sub> and *2*)-related replicons (J. Sun, S. Xiaobo, T. Mikalsen, J. U. Ericson Sollid, A. Sundsfjord, unpublished observations). Other comprehensive studies include vancomycin-resistant *E. faecium* and *E. faecalis* strains causing hospital outbreaks in five continents, from 1986 to date (A. R. Freitas, M. V. Francia, L. Peixe, C. Novais, L. B. Jensen, R. J. Willems, F. Baquero, T. M. Coque, unpublished observations). Among *E. faecium*, mostly CC17-related, a high diversity of *rep* types could be identified; small [pB82 (*rep*<sub>11</sub>), pRII (*rep*<sub>14</sub>), pEF418 (*rep*<sub>18</sub>), pCIZ2 (*rep*<sub>unique</sub>)] or medium to large plasmids [Inc18 (*rep*<sub>1</sub> and *2*), pRUM (*rep*<sub>17</sub>), pHTβ (*rep*<sub>unique</sub>)], with *vanA* linked to Inc18- and pRUM-like plasmids in most cases. Vancomycin-resistant *E. faecalis* isolates belonging to major clonal complexes (CC2, CC9 and CC87) contained a lower diversity of plasmids, which were mostly associated with the narrow host pheromone-responsive pADI (*rep*<sub>9</sub>) and Inc18-type (*rep*<sub>1</sub> and *2*) plasmids.

Linkage of clinically important resistance determinants to specific replicon types in enterococci is of interest for predicting potential transfer to other bacterial genera by conjugative broad host range plasmids. The application of pulsed-field gel electrophoresis of S1-nuclease-digested enterococcal DNA has proved very useful for the identification and sizing of enterococcal plasmids because they appear as linearized bands (5–400 kb) on a faint genomic background [25–27]. Physical linkage between defined plasmid *rep* types and resistance determinants can be visualized by co-hybridization analysis of linearized plasmid DNAs [25]. Co-hybridization

analysis and plasmid sequencing have shown linkage of the *vanA* operon to Inc18-, pHTbeta- and pRUM-related plasmids in *E. faecium* [25,28–31].

Plasmid genomic analysis in enterococci has been hindered by the comparably high number of extrachromosomal elements in many (especially clinical) strains; the tremendous size of several of these plasmids and the multicopy insertion sequence (IS) elements located on them complicating raw data and contig assembly. Sequencing of large multiresistance megaplasmids (>200 kb) are in progress, revealing the presence of multiple resistance determinants linked to known virulence-associated proteins (J. A. Laverde-Gómez, G. Werner, unpublished observations).

## Transposable Elements Involved in Spread of Resistance in Enterococci

IS elements are the simplest transposable elements encoding only the enzyme(s) necessary for their own transposition. Transposons can be classified into three groups: (i) composite transposons; (ii) Tn3 family transposons; and (iii) conjugative transposons [32]. A schematic presentation of enterococcal transposons from each of these groups is presented in Fig. 1. An extensive list of transposons containing specific resistance and virulence determinants described in enterococci is given in Table 1.

Composite transposons owe their intracellular mobility to the presence of flanking copies of IS of the same family that act together to move the DNA between them and have mostly been associated with high-level gentamicin resistance (HLGR) or *vanB1*-related glycopeptide resistance (Table 1). The Tn3 family of transposons move (transpose) intracellularly within or between different replicons through a replicative

mechanism promoted by a transposase (TnpA) and a resolvase (TnpR) [33]. Tn3-like transposons mediate high-level glycopeptide (VanA-type) and macrolide-lincosamide-streptogramin B (MLS<sub>B</sub>) resistance in enterococci (Table 1). Conjugative transposons (CTns) have been classified as integrative conjugative elements (ICE) that encode all information necessary for their own excision, conjugation and integration into a new host [34]. CTns have been associated with resistance to tetracyclines and MLS-antibiotics as well as *vanB2*-related glycopeptide resistance (Table 1). Members of the Tn916/Tn1545-family are considered to comprise classical conjugative transposons with a broad host range carrying clinically relevant resistance determinants between a larger diversity of bacterial genera (Table 1) [35,36]. Importantly, CTns are also able to co-transfer other plasmids and transposons, as well as facilitate transfer of large chromosomal fragments between strains [32].

Whole-genome sequencing and comparative genomic analysis have given insights into the mobile enterococcal DNA pool [38–40]. The IS appear to represent the major part of MGEs in clinical enterococcal isolates, and frequently found IS-families include IS3, IS6, IS30, IS256 and ISL3 [40–57]. Other IS element families found include IS4, IS66, IS110, IS200/IS605, IS982, IS1182 and IS1380 [40–42,49,51,58–60]. IS16 is prevalent in hospital-adapted subpopulations of *E. faecium*, but has also been described in clinical *E. faecalis* strains and as part of pRUM-like plasmids [40–43].

We have limited knowledge concerning the prevalence and distribution of defined composite transposons. This is a result of their highly complex, dynamic and modular structure, which does not allow simplistic detection and typing methods. Recently, the ubiquitous presence of Tn916 and Tn1549/5382-like elements has been demonstrated in oral and faecal human metagenomes from several countries using

**FIG. 1.** Schematic presentation of transposons from each of the three transposon groups transferring resistance genes in enterococci. *aac6'-aph2''*, aminoglycoside-6'-N-acetyltransferase-2''-O-phosphoryltransferase; *tnpA*, transposase; *tnpR*, resolvase; *vanA/vanB2* cluster, clusters conferring vancomycin resistance; *tra* region, region containing transfer genes; *oriT*, origin of transfer; *xis*, excisionase; *int*, integrase.


### Tn5281 composite transposon



### Tn1546 transposon of the Tn3 family



### Tn1549/Tn5382-like conjugative transposon



**TABLE 1.** Transposons containing antimicrobial resistance or virulence genes found in enterococci

Tn3 transposon family					
Tn designation(s)	Size (kb)	Replication and resolution module	Resistance (gene)	Host range <sup>a</sup>	Reference
Tn917 (Tn551) Tn3871	5.3 5.1	TnpA transposase ( <i>tnpA</i> ) and TnpR resolvase ( <i>tnpR</i> )	Macrolides, lincosamides and streptogramin B ( <i>ermB</i> )	<i>Bacillus</i> , <i>Enterococcus</i> , <i>Lactococcus</i> , <i>Listeria</i> , <i>Paenibacillus</i> , <i>Staphylococcus</i> , <i>Streptococcus</i>	[157–159]
Tn1546 and variants	11	TnpA transposase ( <i>tnpA</i> ) and TnpR resolvase ( <i>tnpR</i> )	Glycopeptides ( <i>vanA</i> )	<i>Enterococcus</i> , <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Bacillus circulans</i> , <i>Oerskovia turbata</i> , <i>Arcanobacterium haemolyticum</i> , <i>Paenibacillus</i> , <i>Rhodococcus</i>	[10,13]
Composite transposons					
Tn designation(s)	Size (kb)	Flanking IS elements	Resistance (gene)	Host range <sup>a</sup>	Reference
Tn5281 (Tn4001), Tn3706 and variants of these	4.7	Inverted copies of IS256 or IS257	Aminoglycosides ( <i>aac(6')</i> - <i>aph(2')</i> )	<i>Enterococcus</i> , <i>Mycoplasma</i> , <i>Staphylococcus</i> , <i>Streptococcus</i>	[124,160–165]
Tn924	27	IS257-like	Aminoglycoside ( <i>aac(6')</i> - <i>aph(2')</i> )	<i>Enterococcus faecalis</i>	[166]
Tn1547	64	IS16 and IS256-like in a direct orientation	Vancomycin ( <i>vanB1</i> )	<i>Enterococcus faecalis</i>	[43]
Tn5384	26	Directly repeated copies of IS256	Aminoglycosides ( <i>aac(6')</i> - <i>aph(2')</i> ), erythromycin ( <i>ermB</i> ), mercuric chloride ( <i>merX</i> )	<i>Enterococcus faecalis</i>	[167,168]
Tn5385	Ca 65	Directly repeated copies of IS1216	Aminoglycosides ( <i>aac(6')</i> - <i>aph(2')</i> ), erythromycin ( <i>ermB</i> ), mercuric chloride ( <i>mer</i> ), streptomycin ( <i>aadE</i> ), tetracycline-minocycline ( <i>tetM</i> ), penicillin ( <i>blaZ</i> )	<i>Enterococcus faecalis</i>	[69,169]
Tn5405 and variants	12	Inverted copies of IS1182	Aminoglycosides ( <i>aphA-3</i> , <i>aadE</i> ), streptothricin (similar to <i>sat4</i> )	<i>Enterococcus faecium</i> , <i>Staphylococcus</i>	[60,170]
Tn5482	26–30	IS1216V	Glycopeptides ( <i>vanA</i> )	<i>Enterococcus faecium</i> , <i>Enterococcus faecalis</i>	[44]
Tn5506	39	IS1216V2 in inverted direction of IS1216V1 with IS1252 insertion	Glycopeptides ( <i>vanA</i> )	<i>Enterococcus faecium</i>	[50]
Conjugative transposons (ICEs)					
Tn designation(s)	Size (kb)	Recombination module	Resistance/virulence (gene)	Host range <sup>a</sup>	Reference
Tn916, Tn918, Tn919, Tn920, Tn925, Tn3702, Tn5031, Tn5032, Tn5033, Tn5381, Tn5383	15–23	Tn916 tyrosine recombinase/integrase ( <i>int</i> ) and excisionase ( <i>xis</i> )	Tetracycline-minocycline ( <i>tetM</i> )	Species within the following genera: <i>Acholeplasma</i> , <i>Acinetobacter</i> , <i>Actinobacillus</i> , <i>Alcaligenes</i> , <i>Bacillus</i> , <i>Butyrivibrio</i> , <i>Citrobacter</i> , <i>Clostridium</i> , <i>Enterococcus</i> , <i>Erysipelothrix</i> , <i>Escherichia</i> , <i>Fusobacterium</i> , <i>Granulicatella</i> , <i>Haemophilus</i> , <i>Lactobacillus</i> , <i>Lactococcus</i> , <i>Leuconostoc</i> , <i>Listeria</i> , <i>Mycoplasma</i> , <i>Neisseria</i> , <i>Peptostreptococcus</i> , <i>Pseudomonas</i> , <i>Staphylococcus</i> , <i>Streptococcus</i> , <i>Thermus</i> , <i>Ureaplasma</i> , <i>Veillonella</i>	[171–179]
Tn1545 and variants	Ca 25	Tyrosine recombinase/ integrase ( <i>int</i> ) and excisionase ( <i>xis</i> ) related to those from Tn916	Tetracycline-minocycline ( <i>tetM</i> ), macrolides, lincosamides and streptogramin B ( <i>ermB</i> ), kanamycin ( <i>aphA-3</i> )	Species within the following genera: <i>Bacillus</i> , <i>Bacteroides</i> , <i>Clostridium</i> , <i>Enterococcus</i> , <i>Escherichia</i> , <i>Eubacterium</i> , <i>Lactococcus</i> , <i>Listeria</i> , <i>Neisseria</i> , <i>Roseburia</i> , <i>Staphylococcus</i> , <i>Streptococcus</i> , <i>Ureaplasma</i>	[180,181]
Tn6009	18	Tyrosine recombinase/ integrase ( <i>int</i> ) and excisionase ( <i>xis</i> ) related to those from Tn916	Tetracycline-minocycline ( <i>tetM</i> ), mercury ( <i>mer</i> )	<i>Enterococcus</i> , <i>Klebsiella</i> , <i>Pseudomonas</i> , <i>Serratia</i> , <i>Streptococcus</i>	[182]

TABLE 1. (Continued)

Conjugative transposons					
Tn designation(s)	Size (kb)	Recombination module	Resistance/virulence (gene)	Host range <sup>a</sup>	Reference
Tn5382, Tn1549 and variants of these	27–34	Tyrosine recombinase/ integrase ( <i>int</i> ) and excisionase ( <i>xis</i> ) related to those from Tn916	Vancomycin ( <i>vanB2</i> )	<i>Enterococcus faecium</i> , <i>Enterococcus faecalis</i> , <i>Enterococcus gallinarum</i> , <i>Enterococcus hirae</i> , <i>Clostridium</i> , <i>Streptococcus</i> , <i>Ruminococcus</i> , <i>Eggerthella lentha</i>	[42,63,64,104, 105,108,183]
Tn5386	29	Tyrosine recombinase/ integrase ( <i>int</i> ) and excisionase ( <i>xis</i> ) related to those from Tn916	Lantibiotics ( <i>spa</i> ), putative surface exposed collagen adhesion protein	<i>Enterococcus faecium</i>	[184]
OGIRF conjugative transposon homologue	Ca 49	Putative phage-related integrase, lacks excisionase homologue	Putative adhesin proteins	<i>Enterococcus faecalis</i>	[39]
Tn6000/EfcTnI	33	Excisionase and integrase related to those from <i>S. aureus</i> pathogenicity islands SaPIbov and SaPIbov2	Tetracycline ( <i>tetS</i> )	<i>Enterococcus faecalis</i> , <i>Enterococcus faecium</i> , <i>Enterococcus casseliflavus</i> , <i>Enterococcus</i> spp.	[36,185; C. Novais, A. Freitas, F. Baquero, L. Peixe, M. Zervos, A. Roberts, T. M. Coque, unpublished observations]
CW459TetM/Tn5801 (Tn6104)	Ca 25	Integrase not related to Tn916	Tetracycline-minocycline ( <i>tetM</i> )	<i>Enterococcus faecium</i> , <i>Enterococcus faecalis</i> , <i>Enterococcus</i> spp., <i>Staphylococcus aureus</i> , <i>Clostridium perfringens</i>	C. Novais, A. Freitas, F. Baquero, L. Peixe, T. M. Coque, unpublished observations
V583 <i>vanB1</i> -containing element	55	TndX-like large serine recombinase	Vancomycin ( <i>vanB1</i> )	<i>Enterococcus faecalis</i>	[38]
Tn5397 and variants	Ca 21	TndX-like large serine recombinase	Tetracycline-minocycline ( <i>tetM</i> )	<i>Enterococcus faecium</i> , <i>Enterococcus hirae</i> , <i>Clostridium difficile</i> , <i>Bacillus subtilis</i> , <i>Streptococcus</i>	[186,187]
Tn950	47	Unknown, not homologous with Tn916	Erythromycin	<i>Enterococcus faecium</i>	[188]

<sup>a</sup>Host range includes both hosts into which the elements have been introduced and hosts in which they occur naturally.

integrase-specific probes for detection [61,62]. Detection of the most common CTNs associated with tetracycline, macrolide or chloramphenicol resistance revealed the wide spread of sequences related to Tn916/Tn1545, Tn5397, and also CW459TetM/Tn5801 and CTn6000 among enterococci from different sources (C. Novais, A. Freitas, F. Baquero, L. Peixe, M. Zervos, A. Roberts, T. M. Coque, unpublished observations), while Tn1549/5382-like elements are the most common vehicle for dissemination of VanB-type glycopeptide resistance in enterococci [42,63–66].

### Transposable Elements: Contribution to Genome Plasticity and Phenotypic Variation

Genomic comparisons of host-adapted and less host-adapted species suggest a higher prevalence of mobile elements among the latter, specifying a comparably high genomic stability in very well host-adapted species and a comparably high genomic plasticity in variants with changing hosts or approaching new ecological niches [67]. The latter may be true for the recently emerging hospital-adapted clonal types

of *E. faecium* and *E. faecalis* and may explain their large content of IS and other mobile elements among their genomes compared to other representatives, also within the same species (*E. faecalis* OGIRF/Symbioflor 1) [39,68]. Transposable elements may contribute to genome plasticity by several mechanisms. Multiple copies of related transposable elements are substrates for homologous recombination within and between different DNA elements mediating rearrangements in chromosomal and plasmid DNA [50,69,70]. Moreover, transposable elements may alter the expression of genes either through disruption of genes by insertion within the coding region or by integrations in the promoter region, either disrupting the existing promoter or providing a new and more efficient promoter. Enterococcal ISs found to contribute to the formation of functional promoters include IS3, IS30, IS256 and IS982 [71]. IS1542 and IS1191, both belonging the IS256 family, have been shown to contribute to constitutive expression of the *vanA* operon and to a higher level of biofilm formation in a virulent *E. faecalis*, respectively [45,47]. Moreover, the constitutive expression of vancomycin resistance in *vanD* isolates can be a result of ISEfa4 (IS200/IS605 family) insertion in the *vanD* operon regulator component determinant *vanS<sub>D</sub>* [59] and IS19 (IS982 family), disrupt-

ing the *ddl* gene encoding the housekeeping D-Alanine-D-Alanine ligase. Absence of D,D-carboxypeptidase activity in an *E. faecium* strain occurred as a result of *ISEfa9* (IS3 family) insertion in *vanY<sub>D</sub>* which contributes to lowering the glycopeptide resistance level [49].

## MGE and their Contribution to Specific Resistance Phenotypes

### Transferable resistance to vancomycin

Subsequent to its first description [8,9], the vancomycin resistance operon (*van*) alphabet has expanded to eight in enterococci, namely the acquired *vanA*, *vanB*, *vanD*, *vanE*, *vanG*, *vanL* [72] and *vanN* [73] in addition to the intrinsic *vanC* genotype in *E. gallinarum* and *E. casseliflavus*. Their characteristics and species distribution are summarized in Table 2. The *vanA* and *vanB* genotypes are the most commonly encountered forms of acquired glycopeptide resistance and have primarily been reported in *E. faecium* and *E. faecalis*. The mechanisms of resistance involve modification of the vancomycin-binding target by synthesis of peptidoglycan precursors with peptide sidechains that terminate in D-lactate (*vanA*, *vanB* and *vanD*) or D-serine (*vanC*, *vanE*,

*vanG*, *vanL* and probably *vanN*) for which vancomycin has lower affinity than for the normal D-alanine sidechain terminus [74–77]. Detailed descriptions are provided in recent reviews [78,79].

The *vanA* gene cluster is carried by Tn1546, a Tn3 derivative that is most often found as a part of nonconjugative or conjugative plasmids [10,44,50,80]. Broad host range Inc18 plasmids have been associated with transfer of Tn1546 from enterococci to *Staphylococcus aureus* [81]. The *vanB* operon can be transferred between enterococci as part of large conjugative chromosomal elements or by conjugative plasmids [42,43,63,64,82–89]. The *vanB* ligase gene has been divided into three subtypes, *vanB1–3*, based on nucleotide sequence differences [90–92]. The *vanB2* subtype as an integral part of Tn1549/Tn5382-like conjugative transposons [42,63–66] is the most widespread *vanB* type in clinical enterococci [42,65,80,89,93–100]. Interestingly, the first description of a *vanB2*-Tn1549-like element in pheromone-responsive (pCF10-like) plasmids in *E. faecalis* strains was recently reported from Japan [89]. The occurrence of *vanB1* has only been described for single isolates as an integral part of putative ICE or composite transposons [38,43].

A high prevalence of the *vanB2* gene has been observed in community and hospital human faecal specimens in the

**TABLE 2.** Glycopeptide resistance due to *van*-type gene clusters

Resistance level	Acquired							Intrinsic
	High	Variable	Moderate	Low			Low	
Type	VanA	VanB	VanD	VanE	VanG	VanL	VanN	VanC
MIC in mg/L:								
Vancomycin	≥16	≥4	≥64	6–32	12–16	8	8	2–32
Teicoplanin	>8	0.5–1	4–64	0.5	0.5			0.5–1
Expression	Inducible	Inducible	Constitutive/ Inducible ( <i>vanD2</i> )	Inducible/ (Constitutive)	Inducible			Constitutive/ Inducible
van ligase gene	<i>vanA</i>	<i>vanB1–B3</i>	<i>vanD1–5</i>	<i>vanE</i>	<i>vanG1–2</i>	<i>vanL</i>	<i>vanN</i>	<i>vanC1–C3</i>
Modified target	D-alanine-D-lactate	D-alanine-D-lactate	D-alanine-D-lactate	D-alanine-D-serine	D-alanine-D-serine	D-alanine-D-serine	D-alanine-D-serine?	D-alanine-D-serine
Conjugative transfer	Yes	Yes	No	No	Yes	No	Yes	No
Location	Plasmid/chromosome on transposon(s)	Plasmid/chromosome ± transposon/ICE <sup>a</sup>	Chromosome	Chromosome	Chromosome on possible ICE	Chromosome?	Plasmid	Chromosome
Distribution	<i>Enterococcus faecium</i> <i>Enterococcus faecalis</i> <i>Enterococcus avium</i> <i>Enterococcus casseliflavus</i> <i>Enterococcus durans</i> <i>Enterococcus gallinarum</i> <i>Enterococcus hirae</i> <i>Enterococcus mundtii</i> <i>Enterococcus raffinosus</i> <i>Staphylococcus aureus</i> <i>Bacillus circulans</i> <i>Oerskovia turbata</i> <i>Arcanobacterium haemolyticum</i> <i>Paenibacillus</i> <i>Rhodococcus</i>	<i>Enterococcus faecium</i> <i>Enterococcus faecalis</i> <i>Enterococcus casseliflavus</i> <i>Enterococcus durans</i> <i>Enterococcus gallinarum</i> <i>Enterococcus hirae</i> <i>Staphylococcus epidermidis</i> <i>Streptococcus</i> <i>Clostridium</i> <i>Ruminococcus</i> <i>Eggerthella</i>	<i>Enterococcus faecium</i> <i>Enterococcus faecalis</i> <i>Enterococcus avium</i> <i>Enterococcus gallinarum</i> <i>Enterococcus raffinosus</i> Non-enterococcal faecal flora	<i>Enterococcus faecalis</i>	<i>Enterococcus faecalis</i> Non-enterococcal faecal flora	<i>Enterococcus faecalis</i>	<i>Enterococcus faecium</i>	<i>Enterococcus gallinarum</i> – <i>vanC1</i> <i>Enterococcus casseliflavus</i> – <i>vanC2/3</i>

<sup>a</sup>ICE, integrative conjugative element.



absence of cultivable vancomycin-resistant enterococci (VRE) [101–103]. This could be explained by the presence of *vanB2-Tn1549/Tn5382*-like elements identified in other bacterial genera belonging to the normal intestinal flora, such as *Clostridium*, *Ruminococcus*, *Eggerthella* and *Streptococcus* [104–106]. The high rates of non-enterococcal *vanB* in faecal samples result in a low predictive positive value for VRE using PCR-detection of *vanB* directly from faecal samples [103], whereas detection of *vanA* is considered highly specific [107]. These observations are also consistent with the recent description of *Tn1549* integrase in five of six faecal metagenomes from several European countries [62]. Furthermore, conjugative transfer of *vanB2-Tn1549/Tn5382*-like elements from *Clostridium symbiosum* to enterococci in the gut of gnotobiotic mice was recently described during vancomycin selection [108]. Thus, *vanB2* genes are already part of the normal faecal flora, and treatment with vancomycin might select for accidental transfer events to relevant human Gram-positive pathogens including enterococci. Indeed, rates of infections with *vanB*-type VRE (mainly *E. faecium*) are increasing in several European countries [109]. Most worrying is the finding of *vanA*-type vancomycin resistance in methicillin-resistant *S. aureus* (MRSA): VRSA. A detailed description of VRSA is provided elsewhere [81]. *S. aureus* may coexist with VRE in the gastrointestinal tract or in superficial wounds, providing a likely reservoir for development of VRSA [110]. Most recent outbreaks and clinical infections with VRE worldwide are a result of genetic lineages of hospital-adapted *E. faecium* [57,96,100,111–116] that have acquired several virulence and resistance determinants, increasing its ability to survive and cause disease in nosocomial environments [117–122].

#### High-level resistance to aminoglycosides

Acquisition of HLGR abolishes the synergistic bactericidal effect in combination with cell wall active agents in the treatment of severe enterococcal infections. The HLGR phenotype is most commonly associated with the bifunctional enzyme AAC(6′)-Ie-APH(2′′)-Ia and the APH(2′′)-I-family. The composite transposon *Tn5281* (IS256-related) or its variants have been shown to harbor the *aac(6′)-Ie-aph(2′′)* determinant as part of conjugative enterococcal plasmids [123,124]. pMG1 is a fully sequenced conjugative pHTbeta-like 65-kb plasmid from *E. faecium* containing a *Tn5281*-like element [31]. pMG1-related replicons have been associated with transferable VanA-type vancomycin resistance and HLGR in *E. faecium* in the USA, Japan, the UK and the Netherlands [125,126]. HLGR in invasive *E. faecium* and *E. faecalis* strains in European countries is increasing [127,128; <http://www.rivm.nl/earss/>].

#### Linezolid resistance

Linezolid is an oxazolidinone antibiotic used in the treatment of infections caused by multidrug-resistant enterococci. Resistance, still uncommon in enterococci, is associated with mutations in central loop domain V of the 23S ribosomal RNA gene that affect the binding affinity between target and drug [129]. The first mutation in a single 23S allele is the key event in the development of linezolid resistance. Exchange of alleles via recombination occurs then readily, leading to a stepwise increasing resistance level [130]. If not all alleles have recombined, the resistance genotype is reversible. This type of acquired resistance is not transferable and spreads only clonally. Importantly, a new form of linezolid resistance has recently been described in staphylococci, Cfr, for chloramphenicol-florfenicol resistance [131]. The *cfr* determinant was originally described as plasmid-mediated and encodes a methylase that modifies the 23S rRNA ribosomal subunit, conferring cross-resistance to linezolid. The first *cfr* determinant was present in a genetic context including *ermB* and it was suggested to be of enterococcal origin. The *cfr* determinant has so far not been described in enterococci.

#### Resistance to quinopristin-dalfopristin (Q/D)

*E. faecalis* is intrinsically resistant to Q/D as a result of the presence of the *lsa* determinant, whose function remains to be established [132]. Several mechanisms of acquired Q/D resistance have been described in *E. faecium*, including 23S rRNA gene methylation, enzymatic inactivation of streptogramin A and efflux pumps [133,134]. Enterococcal plasmids harbouring *vat(D)* and *vat(E)* determinants encoding acetyltransferases and determinants for MLS<sub>B</sub> resistance inactivating streptogramins A and B have been described [135–137].

#### Resistance to tigecycline

Only sporadic enterococcal isolates with phenotypic intermediate or borderline resistance (MIC = 0.5/≥1 mg/L) are reported in larger clinical trials. The molecular basis of tigecycline resistance in a single clinical *E. faecalis* isolate from an intensive care unit patient after prolonged tigecycline therapy has been investigated. Tigecycline resistance was stable in this epidemic, hospital-adapted strain type (ST6/clonal complex CC2). The mechanism of tigecycline resistance was not clarified [138].

#### Interactions between transferable antimicrobial resistance and pathogenicity

Several enterococcal plasmids have been shown to contain determinants associated with increased capacity for colonization and infection. In some instances, resistance genes and virulence determinants have been physically linked to the

same plasmid. The pheromone-responsive plasmid pADI encodes cytotoxin [139] and aggregation substance (AS) [140]. pCF10 encodes AS [141], and the bacteriocin 2I gene (*bac2I*) is found on pPD1 [142], *bac32* on pTII [143], *bac4I* on pYII4 [144] and *bac43* on pDTI [145]. The *bac4I* gene was also recently described as genetically linked to *vanB2* on the pheromone-responsive and highly conjugative plasmid pMG2200 [89]. AS has been associated with *vanA*-containing pheromone-responsive plasmids in *E. faecalis* [146] and *E. faecium* [50,147]. Furthermore, large conjugative *hylEfm*<sup>+</sup> positive plasmids encoding VanA-type vancomycin resistance and/or resistance to erythromycin and high levels of gentamicin were recently found in *E. faecium* CC17-related strains [27,37]. These plasmids might also have a role in the success of the CC17 *E. faecium* genotype.


#### Persistence of R plasmids in the absence of antibiotic selection

Plasmids have systems that ensure stable inheritance. Interestingly, some R-plasmids contain plasmid addiction systems encoding labile anti-toxins and stable and potent toxins that are able to kill daughter cells that do not inherit the plasmid during cell division (Fig. 2). Thus, once acquired, the R-plasmids ensure their own maintenance even in the absence of antibiotic selection. Three functional plasmid addiction (TA; toxin–antitoxin) systems have been identified in enterococci; Axe-Txe in pRUM [41], epsilon-zeta in pVEF2 [28,148] and par/Fst system in pADI [149]. A prevalent genetic linkage between the pRUM-replicon type and *axe-txe* has recently been demonstrated in *E. faecium* strains, suggesting a successful plasmid backbone combination [25]. The stability of *vanA*-Inc18 plasmids in enterococci in the absence of glyco-

peptide selection has previously been linked to the epsilon-zeta TA-system [29,150]. The results of competitive fitness experiments of *vanA* plasmid-containing vs. recipient *E. faecium* cells revealed that environmental adaptation, *in vivo* gene transfer and plasmid maintenance system(s) favoured long-term VRE persistence without antibiotic selection and compensate for the biological costs of possessing the resistance genes [151]. These earlier *in vitro* experiments accompany recent prevalence studies of VRE among animal reservoirs in the absence of a direct antibiotic selective pressure [152,153].

#### New protection mechanisms against foreign DNA – means to limit the spread of antibiotic resistance?

Clustered, regularly interspaced, short palindromic repeats (CRISPR) have been shown to provide immunity to foreign DNA (phages and plasmids) by direct sequence match [154]. Many CRISPR elements contain phage sequences that provide the host with viral resistance [155,156]. In addition, sequences homologous to the nickase gene in conjugative plasmids of methicillin- and vancomycin-resistant *S. aureus* (MRSA and VRSA) are also found in functional CRISPR elements, interfering with plasmid conjugation and transformation in *Staphylococcus epidermidis* [156]. By contrast, a BLAST search of seven fully sequenced *E. faecium* strains found no functional CRISPR elements, whereas prophage sequences were common (Van Schaik, personal communication). Almost no mobile genetic elements were found in the fully sequenced *E. faecalis* OG1RF that contains two CRISPR elements [39]. An association was noted between the presence of CRISPR and the absence of prophages in this particular strain in contrast to V583. The role of CRISPR in the pro-



**FIG. 2.** Toxin–antitoxin (TA) pairs contribute efficiently to plasmid maintenance in a bacterial population. If TA-carrying plasmids are not inherited by the daughter cells after cell division, the less stable antitoxin is degraded by proteases and the toxin is left free to interact with its intracellular target, which leads to growth inhibition or cell death.



tection of foreign DNA in enterococci remains to be examined.

## Concluding Remarks

Recent progress in enterococcal genome information and in the development of simplistic typing methods has allowed new insights into the prevalence, distribution and genetic content of MGEs in enterococci. This may allow an assessment of their relative contributions to the spread of defined resistance phenotypes and the potential risk for transfer to other bacterial genera by broad host range conjugative elements. The identification of mechanisms involved in maintenance of R-plasmids has provided information that, at least in part, can explain the persistence of defined transferable resistances in the absence of antibiotic selection. The potential use of genetically manipulated CRISPR interference mechanisms to limit the spread of antibiotic resistant enterococci needs to be examined.

## Acknowledgements

We thank T. Z. Pedersen for providing Fig. 2.

## Transparency Declaration

This work was supported by research grants from the Norwegian Research Council (projects no. 165997 and 183653/S10), Northern Norway Regional Health Authority Medical Research Program and the European Commission (LSHE-CT-2007-03410 'ACE'). The authors state that they have no conflict of interest.

## References

1. Hidron AI, Edwards JR, Patel J et al. NHSN annual update: antimicrobial-resistant pathogens associated with healthcare-associated infections: annual summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2006–2007. *Infect Control Hosp Epidemiol* 2008; 29: 996–1011.
2. McDonnell G, Russell AD. Antiseptics and disinfectants: activity, action, and resistance. *Clin Microbiol Rev* 1999; 12: 147–179.
3. Murray BE. The life and times of the *Enterococcus*. *Clin Microbiol Rev* 1990; 3: 46–65.
4. Rice LB. Emergence of vancomycin-resistant enterococci. *Emerg Infect Dis* 2001; 7: 183–187.
5. Courvalin PM, Carlier C, Chabbert YA. Plasmid-linked tetracycline and erythromycin resistance in group D 'streptococcus'. *Ann Inst Pasteur (Paris)* 1972; 123: 755–759.
6. Clewell DB, Yagi Y, Dunny GM et al. Characterization of three plasmid deoxyribonucleic acid molecules in a strain of *Streptococcus faecalis*: identification of a plasmid determining erythromycin resistance. *J Bacteriol* 1974; 117: 283–289.
7. Jacob AE, Hobbs SJ. Conjugal transfer of plasmid-borne multiple antibiotic resistance in *Streptococcus faecalis* var. *zymogenes*. *J Bacteriol* 1974; 117: 360–372.
8. Uttley AH, Collins CH, Naidoo J et al. Vancomycin-resistant enterococci [letter]. *Lancet* 1988; 1: 57–58.
9. Leclercq R, Derlot E, Duval J et al. Plasmid-mediated resistance to vancomycin and teicoplanin in *Enterococcus faecium*. *N Engl J Med* 1988; 319: 157–161.
10. Arthur M, Molinas C, Depardieu F et al. Characterization of Tn1546, a Tn3-related transposon conferring glycopeptide resistance by synthesis of depsipeptide peptidoglycan precursors in *Enterococcus faecium* BM4147. *J Bacteriol* 1993; 175: 117–127.
11. Coque T. Evolutionary biology of pathogenic enterococci. In: Baquero F, Nombela C, Cassell GH et al., eds, *Evolutionary biology of bacterial and fungal pathogens*. Washington, DC: ASM Press, 2008; 501–521.
12. Engel HW, Soedirman N, Rost JA et al. Transferability of macrolide, lincosamide, and streptogramin resistances between group A, B, and D streptococci, *Streptococcus pneumoniae*, and *Staphylococcus aureus*. *J Bacteriol* 1980; 142: 407–413.
13. Weigel LM, Clewell DB, Gill SR et al. Genetic analysis of a high-level vancomycin-resistant isolate of *Staphylococcus aureus*. *Science* 2003; 302: 1569–1571.
14. Clewell DB, Flannagan SE, Jaworski DD. Unconstrained bacterial promiscuity: the Tn916-Tn1545 family of conjugative transposons. *Trends Microbiol* 1995; 3: 229–236.
15. Dunny GM. The peptide pheromone-inducible conjugation system of *Enterococcus faecalis* plasmid pCF10: cell-cell signalling, gene transfer, complexity and evolution. *Philos Trans R Soc Lond B Biol Sci* 2007; 362: 1185–1193.
16. Thomas CM. Evolution and population genetics of bacterial plasmids. In: Funnal BE, Philips GJ, eds, *Plasmid Biology*. Washington, DC: ASM Press, 2004; 509–528.
17. Novick RP. Plasmid incompatibility. *Microbiol Rev* 1987; 51: 381–395.
18. Carattoli A. Resistance plasmid families in Enterobacteriaceae. *Antimicrob Agents Chemother* 2009; 53: 2227–2238.
19. Flannagan SE, Chow JW, Donabedian SM et al. Plasmid content of a vancomycin-resistant *Enterococcus faecalis* isolate from a patient also colonized by *Staphylococcus aureus* with a VanA phenotype. *Antimicrob Agents Chemother* 2003; 47: 3954–3959.
20. Chang S, Sievert DM, Hageman JC et al. Infection with vancomycin-resistant *Staphylococcus aureus* containing the *vanA* resistance gene. *N Engl J Med* 2003; 348: 1342–1347.
21. Clewell DB, Francia MV. Conjugation in Gram-positive bacteria. In: Funnal BE, Philips GJ, eds, *Plasmid Biology*. Washington, DC: ASM Press, 2004; 227–256.
22. Weaver KE, Kwong SM, Firth N et al. The RepA<sub>N</sub> replicons of Gram-positive bacteria: a family of broadly distributed but narrow host range plasmids. *Plasmid* 2009; 61: 94–109.
23. Francia MV, Varsaki A, Garcillan-Barcia MP et al. A classification scheme for mobilization regions of bacterial plasmids. *FEMS Microbiol Rev* 2004; 28: 79–100.
24. Jensen LB, Garcia-Migura L, Valenzuela AJ et al. A classification system for plasmids from enterococci and other Gram-positive bacteria. *J Microbiol Methods* 2010; 80: 25–43.
25. Rosvoll TC, Pedersen T, Sletvold H et al. PCR-based plasmid typing in *Enterococcus faecium* strains reveals widely distributed pRE25-, pRUM-, pIP501- and pHTbeta-related replicons associated with glycopeptide resistance and stabilizing toxin-antitoxin systems. *FEMS Immunol Med Microbiol* 2010; 58: 254–268.

26. Barton BM, Harding GP, Zuccarelli AJ. A general method for detecting and sizing large plasmids. *Anal Biochem* 1995; 226: 235–240.
27. Arias CA, Panesso D, Singh KV *et al.* Cotransfer of antibiotic resistance genes and a *hylEfm*-containing virulence plasmid in *Enterococcus faecium*. *Antimicrob Agents Chemother* 2009; 53: 4240–4246.
28. Sletvold H, Johnsen PJ, Simonsen GS *et al.* Comparative DNA analysis of two *vanA* plasmids from *Enterococcus faecium* strains isolated from poultry and a poultry farmer in Norway. *Antimicrob Agents Chemother* 2007; 51: 736–739.
29. Sletvold H, Johnsen PJ, Hamre I *et al.* Complete sequence of *Enterococcus faecium* pVEF3 and the detection of an omega-epsilon-zeta toxin-antitoxin module and an ABC transporter. *Plasmid* 2008; 60: 75–85.
30. Zhu W, Clark NC, McDougal LK *et al.* Vancomycin-resistant *Staphylococcus aureus* isolates associated with Inc18-like *vanA* plasmids in Michigan. *Antimicrob Agents Chemother* 2008; 52: 452–457.
31. Tanimoto K, Ike Y. Complete nucleotide sequencing and analysis of the 65-kb highly conjugative *Enterococcus faecium* plasmid pMG1: identification of the transfer-related region and the minimum region required for replication. *FEMS Microbiol Lett* 2008; 288: 186–195.
32. Weaver KE, Rice LB, Churchward G. Plasmids and transposons. In: Gilmore MS, ed. *The enterococci: pathogenesis, molecular biology, and antibiotic resistance*. Washington, DC: ASM Press, 2002; 219–264.
33. Chandler M, Mahillon J. Insertion sequences revisited. In: Craig NL, Craigie R, Gellert M *et al.*, eds. *Mobile DNA II*. Washington, DC: ASM Press, 2002; 305–366.
34. Burrus V, Pavlovic G, Decaris B *et al.* Conjugative transposons: the tip of the iceberg. *Mol Microbiol* 2002; 46: 601–610.
35. Flannagan SE, Zitzow LA, Su YA *et al.* Nucleotide sequence of the 18-kb conjugative transposon Tn916 from *Enterococcus faecalis*. *Plasmid* 1994; 32: 350–354.
36. Roberts AP, Mullany P. A modular master on the move: the Tn916 family of mobile genetic elements. *Trends Microbiol* 2009; 17: 251–258.
37. Freitas AR, Tedim AP, Novais C *et al.* Global spread of the colonization-virulence *hylEfm* gene in megaplasms of CC17 *Enterococcus faecium* polyclonal sub-cluster. *Antimicrob Agents Chemother* 2010; doi: 10.1128/AAC.00134-10 [Epub ahead of print].
38. Paulsen IT, Banerjee L, Myers GS *et al.* Role of mobile DNA in the evolution of vancomycin-resistant *Enterococcus faecalis*. *Science* 2003; 299: 2071–2074.
39. Bourgogne A, Garsin DA, Qin X *et al.* Large scale variation in *Enterococcus faecalis* illustrated by the genome analysis of strain OGIRF. *Genome Biol* 2008; 9: R110.
40. Leavis HL, Willems RJ, Van Wamel WJ *et al.* Insertion sequence-driven diversification creates a globally dispersed emerging multiresistant subspecies of *E. faecium*. *PLoS Pathog* 2007; 3: e7.
41. Grady R, Hayes F. Axe-Txe, a broad-spectrum proteic toxin-antitoxin system specified by a multidrug-resistant, clinical isolate of *Enterococcus faecium*. *Mol Microbiol* 2003; 47: 1419–1432.
42. Dahl KH, Lundblad EW, Røkenes TP *et al.* Genetic linkage of the *vanB2* gene cluster to Tn5382 in vancomycin-resistant enterococci and characterization of two novel insertion sequences. *Microbiology* 2000; 146: 1469–1479.
43. Quintiliani R Jr, Courvalin P. Characterization of Tn1547, a composite transposon flanked by the IS16 and IS256-like elements, that confers vancomycin resistance in *Enterococcus faecalis* BM4281. *Gene* 1996; 172: 1–8.
44. Handwerger S, Skoble J. Identification of chromosomal mobile element conferring high-level vancomycin resistance in *Enterococcus faecium*. *Antimicrob Agents Chemother* 1995; 39: 2446–2453.
45. Brown AR, Townsley AC, Amyes SG. Diversity of Tn1546 elements in clinical isolates of glycopeptide-resistant enterococci from Scottish hospitals. *Antimicrob Agents Chemother* 2001; 45: 1309–1311.
46. Camargo IL, Zanella RC, Brandileone MC *et al.* Occurrence of insertion sequences within the genomes and Tn1546-like elements of glycopeptide-resistant enterococci isolated in Brazil, and identification of a novel element, ISEfa5. *Int J Med Microbiol* 2005; 294: 513–519.
47. Coburn PS, Baghdayan AS, Dolan GT *et al.* An AraC-type transcriptional regulator encoded on the *Enterococcus faecalis* pathogenicity island contributes to pathogenesis and intracellular macrophage survival. *Infect Immun* 2008; 76: 5668–5676.
48. Darini AL, Palepou MF, James D *et al.* Disruption of *vanS* by IS1216V in a clinical isolate of *Enterococcus faecium* with VanA glycopeptide resistance [letter]. *Antimicrob Agents Chemother* 1999; 43: 995–996.
49. Depardieu F, Foucault ML, Bell J *et al.* New combinations of mutations in VanD-Type vancomycin-resistant *Enterococcus faecium*, *Enterococcus faecalis*, and *Enterococcus avium* strains. *Antimicrob Agents Chemother* 2009; 53: 1952–1963.
50. Heaton MP, Discotto LF, Pucci MJ *et al.* Mobilization of vancomycin resistance by transposon-mediated fusion of a VanA plasmid with an *Enterococcus faecium* sex pheromone-response plasmid. *Gene* 1996; 171: 9–17.
51. Huh JY, Lee WG, Lee K *et al.* Distribution of insertion sequences associated with Tn1546-like elements among *Enterococcus faecium* isolates from patients in Korea. *J Clin Microbiol* 2004; 42: 1897–1902.
52. Naas T, Fortineau N, Snanoudj R *et al.* First nosocomial outbreak of vancomycin-resistant *Enterococcus faecium* expressing a VanD-like phenotype associated with a *vanA* genotype. *J Clin Microbiol* 2005; 43: 3642–3649.
53. Novais C, Freitas AR, Sousa JC *et al.* Diversity of Tn1546 and its role in the dissemination of vancomycin-resistant enterococci in Portugal. *Antimicrob Agents Chemother* 2008; 52: 1001–1008.
54. Song JH, Ko KS, Oh WS *et al.* High frequency of vancomycin-resistant *Enterococcus faecium* isolates with VanB phenotype and *vanA* genotype in Korean hospitals. *Diagn Microbiol Infect Dis* 2006; 56: 401–406.
55. Thorisdottir AS, Carias LL, Marshall SH *et al.* IS6770, an enterococcal insertion-like sequence useful for determining the clonal relationship of clinical enterococcal isolates. *J Infect Dis* 1994; 170: 1539–1548.
56. Willems RJ, Top J, Van Den Braak N *et al.* Molecular diversity and evolutionary relationships of Tn1546-like elements in enterococci from humans and animals. *Antimicrob Agents Chemother* 1999; 43: 483–491.
57. Zhu X, Zheng B, Wang S *et al.* Molecular characterisation of outbreak-related strains of vancomycin-resistant *Enterococcus faecium* from an intensive care unit in Beijing, China. *J Hosp Infect* 2009; 72: 147–154.
58. Chen YG, Qu TT, Yu YS *et al.* Insertion sequence ISEcp1-like element connected with a novel *aph(2'')* allele [*aph(2'')*-Ie] conferring high-level gentamicin resistance and a novel streptomycin adenylyl-transferase gene in *Enterococcus*. *J Med Microbiol* 2006; 55: 1521–1525.
59. Depardieu F, Reynolds PE, Courvalin P. VanD-type vancomycin-resistant *Enterococcus faecium* 10/96A. *Antimicrob Agents Chemother* 2003; 47: 7–18.
60. Werner G, Hildebrandt B, Witte W. Linkage of *erm(B)* and *aadE-sat4-aphA-3* in multiple-resistant *Enterococcus faecium* isolates of different ecological origins. *Microb Drug Resist* 2003; 9 (suppl 1): S9–S16.
61. Lancaster H, Bedi R, Wilson M *et al.* The maintenance in the oral cavity of children of tetracycline-resistant bacteria and the genes encoding such resistance. *J Antimicrob Chemother* 2005; 56: 524–531.
62. Seville LA, Patterson AJ, Scott KP *et al.* Distribution of tetracycline and erythromycin resistance genes among human oral and fecal metagenomic DNA. *Microb Drug Resist* 2009; 15: 159–166.
63. Carias LL, Rudin SD, Donskey CJ *et al.* Genetic linkage and cotransfer of a novel, *vanB*-containing transposon (Tn5382) and a low-affinity

- penicillin-binding protein 5 gene in a clinical vancomycin-resistant *Enterococcus faecium* isolate. *J Bacteriol* 1998; 180: 4426–4434.
64. Garnier F, Taourit S, Glaser P et al. Characterization of transposon Tn1549, conferring VanB-type resistance in *Enterococcus* spp. *Microbiology* 2000; 146: 1481–1489.
65. McGregor KF, Nolan C, Young HK et al. Prevalence of the vanB2 gene cluster in vanB glycopeptide-resistant enterococci in the United Kingdom and the Republic of Ireland and its association with a Tn5382-like element. *Antimicrob Agents Chemother* 2001; 45: 367–368.
66. Umeda A, Garnier F, Courvalin P et al. Association between the vanB2 glycopeptide resistance operon and Tn1549 in enterococci from France. *J Antimicrob Chemother* 2002; 50: 253–256.
67. Burrus V, Waldor MK. Shaping bacterial genomes with integrative and conjugative elements. *Res Microbiol* 2004; 155: 376–386.
68. Domann E, Hain T, Ghai R et al. Comparative genomic analysis for the presence of potential enterococcal virulence factors in the probiotic *Enterococcus faecalis* strain Symbioflor 1. *Int J Med Microbiol* 2007; 297: 533–539.
69. Rice LB, Carias LL. Transfer of Tn5385, a composite, multiresistance chromosomal element from *Enterococcus faecalis*. *J Bacteriol* 1998; 180: 714–721.
70. Rice LB, Carias LL, Marshall S et al. Tn5386, a novel Tn916-like mobile element in *Enterococcus faecium* D344R that interacts with Tn916 to yield a large genomic deletion. *J Bacteriol* 2005; 187: 6668–6677.
71. Mahillon J, Chandler M. Insertion sequences. *Microbiol Mol Biol Rev* 1998; 62: 725–774.
72. Boyd DA, Willey BM, Fawcett D et al. Molecular characterization of *Enterococcus faecalis* N06-0364 with low-level vancomycin resistance harboring a novel D-Ala-D-Ser gene cluster, vanL. *Antimicrob Agents Chemother* 2008; 52: 2667–2672.
73. Leclercq R, Lebreton F, Cattior V. Mechanisms of resistance to vancomycin. ESCMID conference on Enterococci: from animal to man 2009; abstract, page 11.
74. Arthur M, Molinas C, Bugg TD et al. Evidence for in vivo incorporation of D-lactate into peptidoglycan precursors of vancomycin-resistant enterococci. *Antimicrob Agents Chemother* 1992; 36: 867–869.
75. Billot Klein D, Gutmann L, Sable S et al. Modification of peptidoglycan precursors is a common feature of the low-level vancomycin-resistant VANB-type *Enterococcus* D366 and of the naturally glycopeptide-resistant species *Lactobacillus casei*, *Pediococcus pentosaceus*, *Leuconostoc mesenteroides*, and *Enterococcus gallinarum*. *J Bacteriol* 1994; 176: 2398–2405.
76. Evers S, Courvalin P. Regulation of VanB-type vancomycin resistance gene expression by the VanS(B)-VanR (B) two-component regulatory system in *Enterococcus faecalis* V583. *J Bacteriol* 1996; 178: 1302–1309.
77. Handwerger S, Pucci MJ, Volk KJ et al. The cytoplasmic peptidoglycan precursor of vancomycin-resistant *Enterococcus faecalis* terminates in lactate. *J Bacteriol* 1992; 174: 5982–5984.
78. Courvalin P. Vancomycin resistance in gram-positive cocci. *Clin Infect Dis* 2006; 42 (suppl 1): S25–S34.
79. Reynolds PE, Courvalin P. Vancomycin resistance in enterococci due to synthesis of precursors terminating in D-alanyl-D-serine. *Antimicrob Agents Chemother* 2005; 49: 21–25.
80. Clark NC, Cooksey RC, Hill BC et al. Characterization of glycopeptide-resistant enterococci from U.S. hospitals. *Antimicrob Agents Chemother* 1993; 37: 2311–2317.
81. Perichon B, Courvalin P. VanA-type vancomycin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2009; 53: 4580–4587.
82. Dahl KH, Røkenes TP, Lundblad EW et al. Nonconjugative transposition of the vanB-containing Tn5382-like element in *Enterococcus faecium*. *Antimicrob Agents Chemother* 2003; 47: 786–789.
83. Hayden MK, Picken RN, Sahm DF. Heterogeneous expression of glycopeptide resistance in enterococci associated with transfer of vanB. *Antimicrob Agents Chemother* 1997; 41: 872–874.
84. Kawalec M, Gniadkowski M, Kedzierska J et al. Selection of a teicoplanin-resistant *Enterococcus faecium* mutant during an outbreak caused by vancomycin-resistant enterococci with the vanB phenotype. *J Clin Microbiol* 2001; 39: 4274–4282.
85. Quintiliani R Jr, Courvalin P. Conjugal transfer of the vancomycin resistance determinant vanB between enterococci involves the movement of large genetic elements from chromosome to chromosome. *FEMS Microbiol Lett* 1994; 119: 359–363.
86. Rice LB, Carias LL, Donskey CL et al. Transferable, plasmid-mediated vanB-type glycopeptide resistance in *Enterococcus faecium*. *Antimicrob Agents Chemother* 1998; 42: 963–964.
87. Woodford N, Jones BL, Baccus Z et al. Linkage of vancomycin and high-level gentamicin resistance genes on the same plasmid in a clinical isolate of *Enterococcus faecalis*. *J Antimicrob Chemother* 1995; 35: 179–184.
88. Woodford N, Morrison D, Johnson AP et al. Plasmid-mediated vanB glycopeptide resistance in enterococci. *Microb Drug Resist* 1995; 1: 235–240.
89. Zheng B, Tomita H, Inoue T et al. Isolation of VanB-type *Enterococcus faecalis* strains from nosocomial infections: first report of the isolation and identification of the pheromone-responsive plasmids pMG2200, encoding VanB-type vancomycin resistance and a Bac41-type bacteriocin, and pMG2201, encoding erythromycin resistance and cytolysin (Hly/Bac). *Antimicrob Agents Chemother* 2009; 53: 735–747.
90. Dahl KH, Simonsen GS, Olsvik Ø et al. Heterogeneity in the vanB gene cluster of genomically diverse clinical strains of vancomycin-resistant enterococci. *Antimicrob Agents Chemother* 1999; 43: 1105–1110.
91. Gold HS, Unal S, Cercenado E et al. A gene conferring resistance to vancomycin but not teicoplanin in isolates of *Enterococcus faecalis* and *Enterococcus faecium* demonstrates homology with vanB, vanA, and vanC genes of enterococci. *Antimicrob Agents Chemother* 1993; 37: 1604–1609.
92. Patel R, Uhl JR, Kohner P et al. DNA sequence variation within vanA, vanB, vanC-1, and vanC-2/3 genes of clinical *Enterococcus* isolates. *Antimicrob Agents Chemother* 1998; 42: 202–205.
93. Demertzi E, Palepou MF, Kaufmann ME et al. Characterisation of VanA and VanB elements from glycopeptide-resistant *Enterococcus faecium* from Greece. *J Med Microbiol* 2001; 50: 682–687.
94. Hanrahan J, Høyen C, Rice LB. Geographic distribution of a large mobile element that transfers ampicillin and vancomycin resistance between *Enterococcus faecium* strains. *Antimicrob Agents Chemother* 2000; 44: 1349–1351.
95. Lee WG, Kim W. Identification of a novel insertion sequence in vanB2-containing *Enterococcus faecium*. *Lett Appl Microbiol* 2003; 36: 186–190.
96. Lopez M, Hormazabal JC, Maldonado A et al. Clonal dissemination of *Enterococcus faecalis* ST201 and *Enterococcus faecium* CC17-ST64 containing Tn5382-vanB2 among 16 hospitals in Chile. *Clin Microbiol Infect* 2009; 15: 586–588.
97. Lorenzo-Diaz F, Delgado T, Reyes-Darias JA et al. Characterization of the first VanB vancomycin-resistant *Enterococcus faecium* isolated in a Spanish hospital. *Curr Microbiol* 2004; 48: 199–203.
98. Lu JJ, Chang TY, Perng CL et al. The vanB2 gene cluster of the majority of vancomycin-resistant *Enterococcus faecium* isolates from Taiwan is associated with the pbp5 gene and is carried by Tn5382 containing a novel insertion sequence. *Antimicrob Agents Chemother* 2005; 49: 3937–3939.
99. Torres C, Escobar S, Portillo A et al. Detection of clonally related vanB2-containing *Enterococcus faecium* strains in two Spanish hospitals. *J Med Microbiol* 2006; 55: 1237–1243.
100. Valdezate S, Labayru C, Navarro A et al. Large clonal outbreak of multidrug-resistant CC17 ST17 *Enterococcus faecium* containing

- Tn5382 in a Spanish hospital. *J Antimicrob Chemother* 2009; 63: 17–20.
101. Domingo MC, Huletsky A, Giroux R *et al*. High prevalence of glycopeptide resistance genes *vanB*, *vanD*, and *vanG* not associated with enterococci in human fecal flora. *Antimicrob Agents Chemother* 2005; 49: 4784–4786.
  102. Graham M, Ballard SA, Grabsch EA *et al*. High rates of fecal carriage of nonenterococcal *vanB* in both children and adults. *Antimicrob Agents Chemother* 2008; 52: 1195–1197.
  103. Young HL, Ballard SA, Roffey P *et al*. Direct detection of *vanB2* using the Roche LightCycler *vanA/B* detection assay to indicate vancomycin-resistant enterococcal carriage - sensitive but not specific. *J Antimicrob Chemother* 2007; 59: 809–810.
  104. Ballard SA, Pertile KK, Lim M *et al*. Molecular characterization of *vanB* elements in naturally occurring gut anaerobes. *Antimicrob Agents Chemother* 2005; 49: 1688–1694.
  105. Dahl KH, Sundsfjord A. Transferable *vanB2* Tn5382-containing elements in fecal streptococcal strains from veal calves. *Antimicrob Agents Chemother* 2003; 47: 2579–2583.
  106. Domingo MC, Huletsky A, Bernal A *et al*. Characterization of a Tn5382-like transposon containing the *vanB2* gene cluster in a *Clostridium* strain isolated from human faeces. *J Antimicrob Chemother* 2005; 55: 466–474.
  107. Stamper PD, Cai M, Lema C *et al*. Comparison of the BD GeneOhm VanR assay to culture for identification of vancomycin-resistant enterococci in rectal and stool specimens. *J Clin Microbiol* 2007; 45: 3360–3365.
  108. Launay A, Ballard SA, Johnson PD *et al*. Transfer of vancomycin resistance transposon Tn1549 from *Clostridium symbiosum* to *Enterococcus* spp. in the gut of gnotobiotic mice. *Antimicrob Agents Chemother* 2006; 50: 1054–1062.
  109. Werner G, Coque TM, Hammerum AM *et al*. Emergence and spread of vancomycin resistance among enterococci in Europe. *Euro Surveill* 2008; 13: 1–11.
  110. Ray AJ, Pultz NJ, Bhalla A *et al*. Coexistence of vancomycin-resistant enterococci and *Staphylococcus aureus* in the intestinal tracts of hospitalized patients. *Clin Infect Dis* 2003; 37: 875–881.
  111. Werner G, Klare I, Fleige C *et al*. Increasing rates of vancomycin resistance among *Enterococcus faecium* isolated from German hospitals between 2004 and 2006 are due to wide clonal dissemination of vancomycin-resistant enterococci and horizontal spread of *vanA* clusters. *Int J Med Microbiol* 2008; 298: 515–527.
  112. Hsieh YC, Lee WS, Ou TY *et al*. Clonal spread of CC17 vancomycin-resistant *Enterococcus faecium* with multilocus sequence type 78 (ST78) and a novel ST444 in Taiwan. *Eur J Clin Microbiol Infect Dis* 2010; 29: 25–30.
  113. Gu L, Cao B, Liu Y *et al*. A new Tn1546 type of VanB phenotype-*vanA* genotype vancomycin-resistant *Enterococcus faecium* isolates in mainland China. *Diagn Microbiol Infect Dis* 2009; 63: 70–75.
  114. Ergani-Ozcan A, Naas T, Baysan BO *et al*. Nosocomial outbreak of vancomycin-resistant *Enterococcus faecium* in a paediatric unit at a Turkish university hospital. *J Antimicrob Chemother* 2008; 61: 1033–1039.
  115. Bonora MG, Olisio D, Lo CG *et al*. Phylogenetic analysis of vancomycin-resistant *Enterococcus faecium* genotypes associated with outbreaks or sporadic infections in Italy. *Microb Drug Resist* 2007; 13: 171–177.
  116. Deplano A, Denis O, Nonhoff C *et al*. Outbreak of hospital-adapted clonal complex-17 vancomycin-resistant *Enterococcus faecium* strain in a haematology unit: role of rapid typing for early control. *J Antimicrob Chemother* 2007; 60: 849–854.
  117. Leavis H, Top J, Shankar N *et al*. A novel putative enterococcal pathogenicity island linked to the *esp* virulence gene of *Enterococcus faecium* and associated with epidemicity. *J Bacteriol* 2004; 186: 672–682.
  118. Leavis HL, Willems RJ, Top J *et al*. High-level ciprofloxacin resistance from point mutations in *gyrA* and *parC* confined to global hospital-adapted clonal lineage CC17 of *Enterococcus faecium*. *J Clin Microbiol* 2006; 44: 1059–1064.
  119. Hendrickx AP, Van Wamel WJ, Posthuma G *et al*. Five genes encoding surface-exposed LPXTG proteins are enriched in hospital-adapted *Enterococcus faecium* clonal complex 17 isolates. *J Bacteriol* 2007; 189: 8321–8332.
  120. Nallapareddy SR, Singh KV, Okhuysen PC *et al*. A functional collagen adhesin gene, *acm*, in clinical isolates of *Enterococcus faecium* correlates with the recent success of this emerging nosocomial pathogen. *Infect Immun* 2008; 76: 4110–4119.
  121. Hendrickx AP, Van Luit-Asbroek M, Schapendonk CM *et al*. SgrA, a nidogen-binding LPXTG surface adhesin implicated in biofilm formation, and EcbA, a collagen binding MSCRAMM, are two novel adhesins of hospital-acquired *Enterococcus faecium*. *Infect Immun* 2009; 77: 5097–5106.
  122. Galloway-Peña JR, Nallapareddy SR, Arias CA *et al*. Analysis of clonality and antibiotic resistance among early clinical isolates of *Enterococcus faecium* in the United States. *J Infect Dis* 2009; 200: 1566–1573.
  123. Watanabe S, Kobayashi N, Quinones D *et al*. Genetic diversity of enterococci harboring the high-level gentamicin resistance gene *aac(6')-Ie-aph(2'')-Ia* or *aph(2'')-Ie* in a Japanese hospital. *Microb Drug Resist* 2009; 15: 185–194.
  124. Simjee S, Manzoor SE, Fraiese AP *et al*. Nature of transposon-mediated high-level gentamicin resistance in *Enterococcus faecalis* isolated in the United Kingdom. *J Antimicrob Chemother* 2000; 45: 565–575.
  125. Tomita H, Pierson C, Lim SK *et al*. Possible connection between a widely disseminated conjugative gentamicin resistance (pMG1-like) plasmid and the emergence of vancomycin resistance in *Enterococcus faecium*. *J Clin Microbiol* 2002; 40: 3326–3333.
  126. Tomita H, Tanimoto K, Hayakawa S *et al*. Highly conjugative pMG1-like plasmids carrying Tn1546-like transposons that encode vancomycin resistance in *Enterococcus faecium*. *J Bacteriol* 2003; 185: 7024–7028.
  127. Lester CH, Sandvang D, Olsen SS *et al*. Emergence of ampicillin-resistant *Enterococcus faecium* in Danish hospitals. *J Antimicrob Chemother* 2008; 62: 1203–1206.
  128. NORM/NORM-VET 2008. Usage of antimicrobial agents and occurrence of antimicrobial resistance in Norway. Tromsø/Oslo: Lundblad Media AS, 2009. ISSN: 1502-2307 (print)/1890-9965 (electronic).
  129. Marshall SH, Donskey CJ, Hutton-Thomas R *et al*. Gene dosage and linezolid resistance in *Enterococcus faecium* and *Enterococcus faecalis*. *Antimicrob Agents Chemother* 2002; 46: 3334–3336.
  130. Willems RJ, Top J, Smith DJ *et al*. Mutations in the DNA mismatch repair proteins MutS and MutL of oxazolidinone-resistant or -susceptible *Enterococcus faecium*. *Antimicrob Agents Chemother* 2003; 47: 3061–3066.
  131. Toh SM, Xiong L, Arias CA *et al*. Acquisition of a natural resistance gene renders a clinical strain of methicillin-resistant *Staphylococcus aureus* resistant to the synthetic antibiotic linezolid. *Mol Microbiol* 2007; 64: 1506–1514.
  132. Singh KV, Murray BE. Differences in the *Enterococcus faecalis* *Isa* locus that influence susceptibility to quinupristin-dalfopristin and clindamycin. *Antimicrob Agents Chemother* 2005; 49: 32–39.
  133. Aksoy DY, Unal S. New antimicrobial agents for the treatment of Gram-positive bacterial infections. *Clin Microbiol Infect* 2008; 14: 411–420.
  134. Werner G, Klare I, Witte W. Molecular analysis of streptogramin resistance in enterococci. *Int J Med Microbiol* 2002; 292: 81–94.
  135. Werner G, Hildebrandt B, Klare I *et al*. Linkage of determinants for streptogramin A, macrolide-lincosamide-streptogramin B, and chloramphenicol resistance on a conjugative plasmid in *Enterococcus*



- faecium* and dissemination of this cluster among streptogramin-resistant enterococci. *Int J Med Microbiol* 2000; 290: 543–548.
136. Werner G, Klare I, Spencker FB et al. Intra-hospital dissemination of quinupristin/dalfopristin- and vancomycin-resistant *Enterococcus faecium* in a paediatric ward of a German hospital. *J Antimicrob Chemother* 2003; 52: 113–115.
137. Simjee S, Zhang Y, Mcdermott PF et al. Heterogeneity of *vat(E)*-carrying plasmids in *Enterococcus faecium* recovered from human and animal sources. *Int J Antimicrob Agents* 2006; 28: 200–205.
138. Werner G, Gfrörer S, Fleige C et al. Tigecycline-resistant *Enterococcus faecalis* strain isolated from a German intensive care unit patient. *J Antimicrob Chemother* 2008; 61: 1182–1183.
139. Clewell DB. Properties of *Enterococcus faecalis* plasmid pAD1, a member of a widely disseminated family of pheromone-responding, conjugative, virulence elements encoding cytolysin. *Plasmid* 2007; 58: 205–227.
140. Galli D, Lottspeich F, Wirth R. Sequence analysis of *Enterococcus faecalis* aggregation substance encoded by the sex pheromone plasmid pAD1. *Mol Microbiol* 1990; 4: 895–904.
141. Olmsted SB, Kao SM, Van Putte LJ et al. Role of the pheromone-inducible surface protein Asc10 in mating aggregate formation and conjugal transfer of the *Enterococcus faecalis* plasmid pCF10. *J Bacteriol* 1991; 173: 7665–7672.
142. Tomita H, Fujimoto S, Tanimoto K et al. Cloning and genetic and sequence analyses of the bacteriocin 21 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pPD1. *J Bacteriol* 1997; 179: 7843–7855.
143. Inoue T, Tomita H, Ike Y. Bac 32, a novel bacteriocin widely disseminated among clinical isolates of *Enterococcus faecium*. *Antimicrob Agents Chemother* 2006; 50: 1202–1212.
144. Tomita H, Kamei E, Ike Y. Cloning and genetic analyses of the bacteriocin 41 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pY114: a novel bacteriocin complemented by two extracellular components (lysin and activator). *J Bacteriol* 2008; 190: 2075–2085.
145. Todokoro D, Tomita H, Inoue T et al. Genetic analysis of bacteriocin 43 of vancomycin-resistant *Enterococcus faecium*. *Appl Environ Microbiol* 2006; 72: 6955–6964.
146. Lim SK, Tanimoto K, Tomita H et al. Pheromone-responsive conjugative vancomycin resistance plasmids in *Enterococcus faecalis* isolates from humans and chicken feces. *Appl Environ Microbiol* 2006; 72: 6544–6553.
147. Magi G, Capretti R, Paoletti C et al. Presence of a *vanA*-carrying pheromone response plasmid (pBRG1) in a clinical isolate of *Enterococcus faecium*. *Antimicrob Agents Chemother* 2003; 47: 1571–1576.
148. Ceglowski P, Boitsov A, Chai S et al. Analysis of the stabilization system of pSM19035-derived plasmid pBT233 in *Bacillus subtilis*. *Gene* 1993; 136: 1–12.
149. Weaver KE, Clewell DB, An F. Identification, characterization, and nucleotide sequence of a region of *Enterococcus faecalis* pheromone-responsive plasmid pAD1 capable of autonomous replication. *J Bacteriol* 1993; 175: 1900–1909.
150. Johnsen PJ, Østerhus JL, Sletvold H et al. Persistence of animal and human glycopeptide-resistant enterococci on two Norwegian poultry farms formerly exposed to avoparcin is associated with a widespread plasmid-mediated *vanA* element within a polyclonal *Enterococcus faecium* population. *Appl Environ Microbiol* 2005; 71: 159–168.
151. Johnsen PJ, Simonsen GS, Olsvik O et al. Stability, persistence, and evolution of plasmid-encoded *VanA* glycopeptide resistance in enterococci in the absence of antibiotic selection *in vitro* and in gnotobiotic mice. *Microb Drug Resist* 2002; 8: 161–170.
152. Nilsson O, Greko C, Top J et al. Spread without known selective pressure of a vancomycin-resistant clone of *Enterococcus faecium* among broilers. *J Antimicrob Chemother* 2009; 63: 868–872.
153. Garcia-Migura L, Liebana E, Jensen LB et al. A longitudinal study to assess the persistence of vancomycin-resistant *Enterococcus faecium* (VREF) on an intensive broiler farm in the United Kingdom. *FEMS Microbiol Lett* 2007; 275: 319–325.
154. Horvath P, Barrangou R. CRISPR/Cas, the immune system of bacteria and archaea. *Science* 2010; 327: 167–170.
155. Barrangou R, Fremaux C, Deveau H et al. CRISPR provides acquired resistance against viruses in prokaryotes. *Science* 2007; 315: 1709–1712.
156. Marraffini LA, Sontheimer EJ. CRISPR interference limits horizontal gene transfer in staphylococci by targeting DNA. *Science* 2008; 19: 322. 1843–5.
157. Tomich PK, An FY, Clewell DB. Properties of erythromycin-inducible transposon Tn917 in *Streptococcus faecalis*. *J Bacteriol* 1980; 141: 1366–1374.
158. Shaw JH, Clewell DB. Complete nucleotide sequence of macrolide-lincosamide-streptogramin B-resistance transposon Tn917 in *Streptococcus faecalis*. *J Bacteriol* 1985; 164: 782–796.
159. Banai M, Leblanc DJ. *Streptococcus faecalis* R plasmid pJH1 contains an erythromycin resistance transposon (Tn3871) similar to transposon Tn917. *J Bacteriol* 1984; 158: 1172–1174.
160. Hodel-Christian SL, Murray BE. Characterization of the gentamicin resistance transposon Tn5281 from *Enterococcus faecalis* and comparison to staphylococcal transposons Tn4001 and Tn4031. *Antimicrob Agents Chemother* 1991; 35: 1147–1152.
161. Simjee S, Fraise AP, Gill MJ. Plasmid heterogeneity and identification of a Tn5281-like element in clinical isolates of high-level gentamicin-resistant *Enterococcus faecium* isolated in the UK. *J Antimicrob Chemother* 1999; 43: 625–635.
162. Straut M, De Cespedes G, Delbos F et al. Molecular typing of *Enterococcus faecalis* strains resistant to high levels of gentamicin and isolated in Romania. *J Antimicrob Chemother* 1997; 39: 483–491.
163. Straut M, De Cespedes G, Horaud T. Plasmid-borne high-level resistance to gentamicin in *Enterococcus hirae*, *Enterococcus avium*, and *Enterococcus raffinosus*. *Antimicrob Agents Chemother* 1996; 40: 1263–1265.
164. Sahm DF, Gilmore MS. High-level gentamicin resistance among enterococci. *Dev Biol Stand* 1995; 85: 99–105.
165. Leelaporn A, Yodkamol K, Waywa D et al. A novel structure of Tn4001-truncated element, type V, in clinical enterococcal isolates and multiplex PCR for detecting aminoglycoside resistance genes. *Int J Antimicrob Agents* 2008; 31: 250–254.
166. Thal LA, Chow JW, Clewell DB et al. Tn924, a chromosome-borne transposon encoding high-level gentamicin resistance in *Enterococcus faecalis*. *Antimicrob Agents Chemother* 1994; 38: 1152–1156.
167. Bonafede ME, Carias LL, Rice LB. Enterococcal transposon Tn5384: evolution of a composite transposon through cointegration of enterococcal and staphylococcal plasmids. *Antimicrob Agents Chemother* 1997; 41: 1854–1858.
168. Rice LB, Marshall SH. Insertions of IS256-like element flanking the chromosomal beta-lactamase gene of *Enterococcus faecalis* CX19. *Antimicrob Agents Chemother* 1994; 38: 693–701.
169. Rice LB. Association of different mobile elements to generate novel integrative elements. *Cell Mol Life Sci* 2002; 59: 2023–2032.
170. Derbise A, De Cespedes G, El SN. Nucleotide sequence of the *Staphylococcus aureus* transposon, Tn5405, carrying aminoglycosides resistance genes. *J Basic Microbiol* 1997; 37: 379–384.
171. Rice LB, Marshall SH, Carias LL. Tn5381, a conjugative transposon identifiable as a circular form in *Enterococcus faecalis*. *J Bacteriol* 1992; 174: 7308–7315.
172. Torres OR, Korman RZ, Zahler SA et al. The conjugative transposon Tn925: enhancement of conjugal transfer by tetracycline in *Enterococcus faecalis* and mobilization of chromosomal genes in *Bacillus subtilis* and *E. faecalis*. *Mol Gen Genet* 1991; 225: 395–400.

173. Franke AE, Clewell DB. Evidence for a chromosome-borne resistance transposon (Tn916) in *Streptococcus faecalis* that is capable of 'conjugal' transfer in the absence of a conjugative plasmid. *J Bacteriol* 1981; 145: 494–502.
174. Fletcher HM, Marri L, Daneo-Moore L. Transposon-916-like elements in clinical isolates of *Enterococcus faecium*. *J Gen Microbiol* 1989; 135: 3067–3077.
175. Clewell DB, An FY, White BA et al. *Streptococcus faecalis* sex pheromone (cAM373) also produced by *Staphylococcus aureus* and identification of a conjugative transposon (Tn918). *J Bacteriol* 1985; 162: 1212–1220.
176. Christie PJ, Korman RZ, Zahler SA et al. Two conjugation systems associated with *Streptococcus faecalis* plasmid pCF10: identification of a conjugative transposon that transfers between *S. faecalis* and *Bacillus subtilis*. *J Bacteriol* 1987; 169: 2529–2536.
177. Casey J, Daly C, Fitzgerald GF. Chromosomal integration of plasmid DNA by homologous recombination in *Enterococcus faecalis* and *Lactococcus lactis* subsp. *lactis* hosts harboring Tn919. *Appl Environ Microbiol* 1991; 57: 2677–2682.
178. Murray BE, An FY, Clewell DB. Plasmids and pheromone response of the beta-lactamase producer *Streptococcus (Enterococcus) faecalis* HH22. *Antimicrob Agents Chemother* 1988; 32: 547–551.
179. Horaud T, Delbos F, De Cespedes G. Tn3702, a conjugative transposon in *Enterococcus faecalis*. *FEMS Microbiol Lett* 1990; 60: 189–194.
180. De Leener E, Martel A, Decostere A et al. Distribution of the *erm(B)* gene, tetracycline resistance genes, and Tn1545-like transposons in macrolide- and lincosamide-resistant enterococci from pigs and humans. *Microb Drug Resist* 2004; 10: 341–345.
181. Poyart-Salmeron C, Trieu-Cuot P, Carlier C et al. Molecular characterization of two proteins involved in the excision of the conjugative transposon Tn1545: homologues with other site-specific recombinases. *EMBO J* 1989; 8: 2425–2433.
182. Soge OO, Beck NK, White TM et al. A novel transposon, Tn6009, composed of a Tn916 element linked with a *Staphylococcus aureus* *mer* operon. *J Antimicrob Chemother* 2008; 62: 674–680.
183. Torres C, Tenorio C, Portillo A et al. Intestinal colonization by *vanA*- or *vanB2*-containing enterococcal isolates of healthy animals in Spain. *Microb Drug Resist* 2003; 9 (suppl 1): S47–S52.
184. Rice LB, Carias LL, Marshall SH et al. Characterization of Tn5386, a Tn916-related mobile element. *Plasmid* 2007; 58: 61–67.
185. Roberts AP, Davis IJ, Seville L et al. Characterization of the ends and target site of a novel tetracycline resistance-encoding conjugative transposon from *Enterococcus faecium* 664.1H1. *J Bacteriol* 2006; 188: 4356–4361.
186. Agersø Y, Pedersen AG, Aarestrup FM. Identification of Tn5397-like and Tn916-like transposons and diversity of the tetracycline resistance gene *tet(M)* in enterococci from humans, pigs and poultry. *J Antimicrob Chemother* 2006; 57: 832–839.
187. Poeta P, Costa D, Igrejas G et al. Phenotypic and genotypic characterization of antimicrobial resistance in faecal enterococci from wild boars (*Sus scrofa*). *Vet Microbiol* 2007; 125: 368–374.
188. Takeuchi K, Tomita H, Fujimoto S et al. Drug resistance of *Enterococcus faecium* clinical isolates and the conjugative transfer of gentamicin and erythromycin resistance traits. *FEMS Microbiol Lett* 2005; 243: 347–354.